

**Supporting Information:**

**Measurement of the Band Bending and Surface Dipole at Chemically Functionalized  
Si(111)/Vacuum Interfaces**

David C. Gleason-Rohrer, Bruce S. Brunschwig,\* and Nathan S. Lewis\*

*Beckman Institute and Kavli Nanoscience Institute*


*Division of Chemistry and Chemical Engineering*

*210 Noyes Laboratory, 127-72*


*California Institute of Technology*

*1200 E. California Blvd.*

*Pasadena, CA 91125*



**Supporting Information: Scheme 1.** Schematic representation of Si(111) surface functionalization.



**Supporting Information: Figure 1.** XPS (a) survey and (b) Br 3d spectra of Si(111) surfaces. Sample surfaces are clean with adventitious carbon and oxygen accounting for 1-3 monolayers on unannealed surfaces. Si(111)-CH<sub>3</sub>[Br] surface concentration of bromine is 10%-30% (10% shown) the concentration of Si(111)-Br surfaces. IR absorbance spectra subtracted from Si(111)-H (c) and (d) demonstrate no loss of characteristic Si(111)-CH<sub>3</sub> peaks upon exposure to bromine. The oxide coverage (not shown) was found to be less than 15%  $\pm$  10% of a monolayer for all samples.